

## Article

# Post-Diagnostic Diet Quality and Mortality in Females with Self-Reported History of Breast or Gynecological Cancers: Results from the Third National Health and Nutrition Examination Survey (NHANES III)

Nena Karavasiloglou <sup>1,2</sup> , Giulia Pestoni <sup>1,2</sup> , David Faeh <sup>1,3</sup>  and Sabine Rohrmann <sup>1,2,\*</sup>

<sup>1</sup> Division of Chronic Disease Epidemiology, Institute for Epidemiology, Biostatistics and Prevention (EBPI), University of Zurich, Hirschengraben 84, CH-8001 Zurich, Switzerland; nena.karavasiloglou@uzh.ch (N.K.); giulia.pestoni@uzh.ch (G.P.); info@davidfaeh.ch (D.F.)

<sup>2</sup> Cancer Registry Zurich and Zug, University Hospital Zurich, Zurich, Switzerland, Vogelsangstrasse 10, CH-8001 Zurich, Switzerland

<sup>3</sup> Health Department—Nutrition and Dietetics, Bern University of Applied Sciences, Falkenplatz 24, CH-3012 Bern, Switzerland

\* Correspondence: sabine.rohrmann@uzh.ch; Tel.: +41-44-63-45256

Received: 11 September 2019; Accepted: 18 October 2019; Published: 23 October 2019



**Abstract:** High quality diets are associated with favorable disease and mortality outcomes in various populations; little and conflicting information is available for female cancer survivors. We investigated the association of post-diagnostic diet quality with mortality in female cancer survivors. Data from 230 women with a previous breast, or gynecological (i.e., ovarian, cervical or uterine) cancer diagnosis in the Third National Health and Nutrition Examination Survey were analyzed. The Healthy Eating Index (HEI) and the Mediterranean Diet Score (MDS) were calculated based on a 24-hour dietary recall interview. Cox proportional hazards regression models were used to calculate multivariable-adjusted hazard ratios (HR) and 95% confidence intervals (CI). Higher HEI score was associated with lower mortality (HR<sub>HEI</sub> total = 0.97, 95% CI: 0.95–0.98, 1 unit increase), but the association for MDS failed to reach statistical significance (HR<sub>MDS</sub> total = 0.87, 95% CI: 0.74–1.04). In subgroup analyses, a statistically significant inverse association was observed between the HEI and mortality; for the MDS, no statistically significant association was apparent. Higher post-diagnostic HEI score was inversely associated with mortality in female cancer survivors, suggesting a protective effect when adhering to the diet captured by the HEI. Additional studies are required in order to investigate underlying mechanisms of the mortality-adherence association.

**Keywords:** dietary patterns; cancer survivors; cohort; adults; mortality

## 1. Introduction

Even though breast cancer and gynecological cancers are the most common cancers in women worldwide [1], little is known about the post-diagnostic habits women adopt and if they are associated with mortality. A recent report indicated that while no recommendations could be made for cancer survivors, there are indications that healthy body weight, physical activity, and dietary factors post-diagnosis may be associated with better survival [2]. This shortage of evidence on the influence of post-diagnostic habits is particularly important since the number of cancer survivors is growing and evidence-based recommendations to prolong survival are needed.

Few studies have reported on the association between post-diagnostic lifestyle and survival in cancer patients and their results are contradicting. Decreased mortality risk in breast cancer survivors who consumed at least five daily portions of fruit and vegetables and were physically active was reported [3]. Older female cancer survivors with higher adherence scores to the 2007 World Cancer Research Fund/American Institute for Cancer Research guidelines on body weight, physical activity, and diet had lower mortality of all causes; lower all-cause mortality was also reported for women who only followed the dietary recommendations [4]. On the other hand, high vegetable, fruit, and dietary fiber intake and low fat consumption was not associated with reduced mortality risk in early stage breast cancer survivors in a different study [5].

The Healthy Eating Index (HEI) is associated with lower risk of chronic diseases and mortality in different populations [6,7]. Higher diet quality (assessed by the HEI) was associated with lower all-cause, cardiovascular disease, and cancer mortality risk in postmenopausal women [8] and a meta-analysis reported similar findings [9]. Additionally, higher post-diagnostic diet quality was associated with lower mortality risk, especially mortality from non-breast cancer causes, in breast cancer survivors [10].

The Mediterranean diet is associated with lower mortality in older populations [11–14], as well as in coronary heart disease [15], cardiovascular disease [16] and diabetic patients [17]. A recent meta-analysis reported no association between adherence to the Mediterranean diet and cancer-specific mortality in cancer patients including data from four studies [18]. Furthermore, no association between the Mediterranean diet with either all-cause mortality or non-breast-cancer-related mortality was reported in breast cancer survivors. However, a statistically significant inverse association with death from non-breast cancer causes was shown in women who reported low physical activity [19].


Based on the possible association of a higher quality post-diagnostic diet with decreased mortality in cancer survivors and the limited number of studies that have reported on it, we believe that the association of post-diagnostic diet quality with mortality in female cancer survivors warrants further investigation.

## 2. Materials and Methods

### 2.1. Population

In this study, data from the Third National Health and Nutrition Examination Survey (NHANES III) were used, due to the long follow-up of the participants. The NHANES III was a nationwide survey, conducted between 1988 and 1994 in the United States. Participants were interviewed and underwent physical examination in a mobile examination center. Detailed description of the methodology of the NHANES III is described elsewhere [20]. NHANES III data are publically available and can be accessed online [21]. Therefore institutional review board approval and oversight were not required for our study.

In this study, only female participants who had self-reported a previous breast, ovarian, cervical or uterine cancer diagnosis were included in the analyses. Participants were defined as cancer survivors if they answered “yes” to the question, “Has a doctor ever told you that you had other cancer?” In following questions participants had to specify the site of their cancer diagnosis and their age at cancer diagnosis (“How old were you when you were first told you had other cancer?”). Participants who also self-reported a skin cancer diagnosis ( $n = 26$ ; answered “yes” to the question, “Has a doctor ever told you that you had skin cancer?”), with missing information on either of the exposure variables ( $n = 4$ ) or any of the confounding variables ( $n = 50$ ) were excluded from the analyses (Figure 1). The final study population included 230 participants. Of these, 110 were breast cancer survivors (BCa) and 120 were gynecological cancer (i.e., ovarian ( $n = 19$ ), cervical ( $n = 54$ ), and uterine cancer ( $n = 47$ ); GynCa) survivors.



**Figure 1.** Flowchart of the study participants.

## 2.2. Outcome Ascertainment

The outcome of interest was all-cause mortality. Probabilistic linkage of the NHANES III with the National Death Index, maintained by the National Center for Health Statistics, was used to obtain mortality information. Detailed description of the linkage was previously described [22]. Previous studies have reported accurate ascertainment of participants' death by the National Death Index [23–25]. Follow-up time was defined as the time from completion of the NHANES III questionnaire (in months) until death from any cause or end of follow-up (December 31, 2011), whichever came first.

## 2.3. Assessment

Dietary information was obtained via a 24-hour dietary recall interview. Data collection was scheduled as such as to include interviews all days of the week and throughout the year. During an interview by trained interviewers, participants were requested to report their food and beverage consumption of the past 24 hours. Additional information including (but not limited to) the quantities, recipes, and place of eating was also recorded. The nutrient values were calculated using recipe ingredient nutrient values found in the US Department of Agriculture's (USDA) Survey Nutrient Database Nutrient Files. The dietary information of the 24-hour dietary recall interview was used to calculate the diet quality indices.

To assess moderate to vigorous physical activity NHANES III participants were asked how frequently (but not for how long) they performed leisure time exercise or physical activities in the past month. Their answers were then transformed to "times per week" using the conversion factor 4.3 weeks per month.

## 2.4. Healthy Eating Index

The HEI, developed by the USDA, includes 10 equally weighted distinct components and the sum of each participant's score in the components constitutes their individual score. Components 1–5 estimate a participant's conformity to the recommendations by the Food Guide Pyramid for the grain, vegetable, fruit, milk, and meat groups. Components 6–9 capture total fat, saturated fat, total cholesterol, and sodium dietary intake. Finally, component 10 estimates the variety in a

participant's diet. The score can range from 0 to 100 and the higher a participant's score on the HEI, the better the diet according to the Dietary Guidelines for Americans and the Food Guide Pyramid [20]. Participants were dichotomized based on their HEI (poor (HEI score < 70); good (HEI score  $\geq$  70)).

### 2.5. Mediterranean Diet Score

The MDS [13] assesses the adherence to the Mediterranean diet by including the consumption of 9 components (legumes, vegetables, fruit and nuts, cereals, fish and seafood, meat and meat products, dairy products, the ratio of monounsaturated to saturated fats and alcohol). The components of the MDS were composed by linking the 24-hour dietary recall data with My Pyramid Equivalents Database for USDA survey food codes. The daily consumption of each of the components (in servings/100 g/day) was calculated for every participant. The median consumption for each component was estimated in our study population and then participants were scored with 0 or 1 for each component based on the relation of their consumption to the study median. Participants with consumption of legumes, vegetables, fruit and nuts, cereals, and fish and seafood above the median received 1 point (per component), whereas consumption of meat and meat products, dairy products and a ratio of monounsaturated to saturated fats lower than the median received 1 point (per component). Finally, participants with alcohol consumption of 5–25 g per day (approximately 0.3 to 2 drinks) received 1 point. The sum of all the points was calculated as each individual's MDS. The MDS can range from 0 to 9. Participants were divided into two groups based on their score; participants with MDS 0–4 were considered to be “non-adherers”, whereas participants with MDS 5–9 were considered to be “adherers”.

### 2.6. Statistical Analysis

Baseline categorical data were expressed as percentages and continuous data as means and standard errors of the mean (SEM). The association between each diet quality index and mortality was assessed using Cox proportional hazards regression models. Regarding confounder adjustment, the first model (Model 1) was adjusted only for age at completion of the NHANES III questionnaire and race/ethnicity (Non-Hispanic white, Non-Hispanic black, Mexican-American, Other). The second model (Model 2) was additionally adjusted for several a priori determined confounders based on the existing literature including: time between cancer diagnosis and completion of the NHANES III questionnaire (continuous, years), socioeconomic status (SES; categories based on the poverty income ratio (PIR), poor/near poor (PIR < 2), middle income ( $2 \leq$  PIR < 4), higher income (PIR  $\geq$  4), or unknown; similar to Suresh et al. [26]), marital status (married/living together, never married/widowed, divorced/separated, or unknown), body mass index (BMI; continuous, kg/m<sup>2</sup>), physical activity (continuous, times of moderate to vigorous physical activity per week), smoking status (never, former, or current smoker), self-reported prevalent chronic diseases at baseline (type 2 diabetes, heart attack, congestive heart failure or stroke; yes, no, or unknown), daily energy intake (continuous, kcal/day) and history of menopausal hormone therapy use (never or no information, ever user). Model 2 was further adjusted for alcohol intake (frequency of beer, wine, and hard liquor intake; times per month) in the analyses for the HEI but not for the MDS, since alcohol intake is already included within this index. The results were presented as hazard ratios (HR) and 95% confidence intervals (CI).

Baseline characteristics were compared between female breast or gynecological cancers survivors and female participants of the NHANES III without a self-reported cancer diagnosis. Sensitivity analysis was performed excluding alcohol consumption from the MDS.

Statistical analyses were performed using SAS (version 9.3, SAS Institute Inc., Cary, NC, USA) software and significance levels were set at  $\alpha = 0.05$ . Sampling weights, adapted according to our total study population and sub-groups, were used in all analyses to account for the complex survey design and survey non-response.

### 3. Results

Overall mean time between cancer diagnosis and completion of the NHANES III questionnaire was 10.4 years (SEM 0.7; breast cancer survivors: 8.6 years, SEM: 0.7; gynecological cancer survivors: 12.0 years, SEM: 1.0). Breast cancer survivors completed the NHANES III questionnaire at an older age and were less likely to be current smokers compared to gynecological cancer survivors (Table 1). Breast cancer survivors were more likely to report higher income and had slightly higher BMI compared to gynecological cancer survivors. Female breast or gynecological cancer survivors included in this project were less physically active and more likely to be current smokers compared to female NHANES III participants who did not report history of cancer (Table 2). Regarding their dietary habits, female breast or gynecological cancer survivors had slightly higher HEI score and increased fruit and vegetable consumption compared to female participants who did not report history of cancer.

**Table 1.** Socioeconomic characteristics and participants' behaviors at completion of the NHANES III questionnaire according to their cancer diagnosis and combined <sup>1</sup>.

	Total Study Population (n = 230)	Breast Cancer Survivors (n = 110)	Gynecological Cancer Survivors (n = 120)
Age at completion of the NHANES III questionnaire (years, mean, SEM)	54.4, 1.5	62.4, 1.6	47.4, 1.6
Age at diagnosis (years, mean, SEM)	44.0, 1.3	53.7, 1.5	35.4, 1.1
Time from the completion of the NHANES III questionnaire until the end of the follow-up (months, mean, SEM)	192.2, 6.9	170.4, 9.6	211.6, 5.7
Race/Ethnicity, %			
Non-Hispanic white	89.8	91.6	88.2
Non-Hispanic black	6.8	5.0	8.4
Mexican-American	2.0	1.5	2.4
Other	1.4	1.9	0.9
Marital status, %			
Married/living together	61.8	60.0	63.3
Never married/widowed	21.0	27.0	15.7
Divorced/separated	17.2	13.0	21.0
Socioeconomic status, %			
Poor or near poor	31.0	25.3	36.0
Middle income	36.6	38.9	34.6
Higher income	25.7	26.9	24.6
Unknown	6.8	8.9	4.8
Smoking status, %			
Never	38.6	42.5	35.1
Former	33.7	40.5	27.7
Current	27.7	16.9	37.3
Moderate and vigorous physical activity (times/week), %			
0	20.8	22.5	19.3
0 to <2	25.8	29.2	22.9
2 to <4	22.4	13.9	29.9
4 to <6	6.5	8.1	5.1
>6	24.4	26.2	22.8
Body mass index (kg/m <sup>2</sup> , mean, SEM)	25.8, 0.5	26.4, 0.5	25.3, 0.5
History of menopausal hormone therapy use, %			
Never or No information	55.7	54.5	56.8
Ever	44.3	45.5	43.2
Prevalent chronic diseases at baseline, %			
Yes	14.3	14.7	14.0
No	83.5	81.1	85.6
Unknown	2.2	4.2	0.4
Healthy Eating Index, %			
Poor (score <70)	60.0	49.7	69.0
Good (score ≥70)	40.0	50.3	31.0
Mediterranean Diet Score, %			
Non-Adherers (score 0–4)	75.7	73.7	77.4
Adherers (score 5–9)	24.3	26.3	22.6

<sup>1</sup> SEM: Standard Error of the Mean.

The association between the diet quality indices and mortality, after an overall mean follow-up time of approximately 16 years, is shown in Table 3. The number of deaths recorded until the end of follow-up was 121. In continuous models, the HEI was statistically significantly associated with lower mortality ( $HR_{HEI\ total} = 0.97$ , 95% CI: 0.95–0.98,  $HR_{HEI\ BCa} = 0.97$ , 95% CI: 0.95–0.99,  $HR_{HEI\ GynCa} = 0.92$ ,

95% CI: 0.89–0.96, in the fully adjusted models, per 1 unit increase in the HEI score). In categorical models, high diet quality (corresponding to HEI  $\geq$  70) was statistically significantly associated with mortality risk in the total study population ( $HR_{HEI \text{ good vs. poor total}} = 0.43$ , 95% CI: 0.29–0.64). Sub-group analyses revealed similar results for gynecological cancer survivors ( $HR_{HEI \text{ good vs. poor GyCa}} = 0.20$ , 95% CI: 0.10–0.43) and breast cancer survivors ( $HR_{HEI \text{ good vs. poor BCa}} = 0.49$ , 95% CI: 0.25–0.97).

**Table 2.** Lifestyle characteristics of females without a self-reported cancer diagnosis vs. females with self-reported breast or gynecological cancers in the NHANES III <sup>1</sup>.

	Females without Self-Reported Cancer (n = 8424)	Females with Self-Reported Breast or Gynecological Cancers (n = 230)
Age at completion of the NHANES III questionnaire (years, mean, SEM)	42.1, 0.5	54.4, 1.5
Race/Ethnicity, %		
Non-Hispanic white	73.8	89.8
Non-Hispanic black	12.6	6.8
Mexican-American	5.2	2.0
Other	8.4	1.4
Marital status, %		
Married/living together	59.2	61.8
Never married/widowed	27.8	21.0
Divorced/separated	12.9	17.2
Unknown	0.1	0.0
Socioeconomic status, %		
Poor	14.3	11.2
Near poor	20.7	19.8
Middle income	35.0	36.6
Higher income	23.2	25.7
Unknown	6.9	6.8
Smoking status, %		
Never	56.4	38.6
Former	18.0	33.7
Current	25.6	27.7
Moderate and vigorous physical activity (times/week), %		
0	18.1	20.8
0 to <2	24.9	25.8
2 to <4	16.9	22.4
4 to <6	7.4	6.5
>6	32.7	24.4
Body mass index (kg/m <sup>2</sup> , mean, SEM)	26.3, 0.2	25.8, 0.5
History of menopausal hormone therapy use, %		
Never	20.8	43.5
Former	7.2	29.3
Current	7.3	15.0
Unknown/No information	64.6	12.2
Breastfeeding, %		
Yes	38.6	44.4
Healthy Eating Index (mean, SEM)	64.2, 0.3	66.3, 1.0
Mediterranean Diet Score (mean, SEM)	3.5, 0.0	3.4, 0.1
Total energy intake (kcal/day, mean, SEM)	1805.2, 15.1	1,741.4, 57.1
Fruit (times/month, mean, SEM)	28.5, 0.6	32.9, 3.2
Vegetables (times/month, mean, SEM)	66.5, 1.0	72.8, 5.5
Red meat (times/month, mean, SEM)	19.7, 0.4	19.8, 1.5

<sup>1</sup> SEM: Standard Error of the Mean.

The MDS was not associated with mortality risk in the total study population ( $HR_{MDS \text{ total}} = 0.87$ , 95% CI: 0.74–1.04, in the fully adjusted model, per 1 unit increase in the MDS). Subgroup analyses revealed similar not statistically significant associations in both groups ( $HR_{MDS \text{ BCa}} = 0.97$ , 95% CI: 0.82–1.16;  $HR_{MDS \text{ GynCa}} = 0.77$ , 95% CI: 0.57–1.04). In categorical models, the categories of the MDS were not statistically significantly associated with mortality risk in the total population ( $HR_{MDS \text{ Adherers vs. Non-Adherers}} = 0.67$ , 95% CI: 0.41–1.11) or the subgroups. Excluding alcohol consumption from the MDS did not materially modify our results (data not shown).

**Table 3.** The association between diet quality indices and mortality for all participants and according to their cancer diagnosis <sup>1</sup>.

	Total Study Population ( <i>n</i> = 230)		Breast Cancer Survivors ( <i>n</i> = 110)		Gynecological Cancer Survivors ( <i>n</i> = 120)	
	Model 1	Model 2	Model 1	Model 2	Model 1	Model 2
Healthy Eating Index	0.97 (0.95–0.99)	0.97 (0.95–0.98)	0.97 (0.95–0.99)	0.97 (0.95–0.99)	0.95 (0.93–0.97)	0.92 (0.89–0.96)
Poor (<70)	1.00	1.00	1.00	1.00	1.00	1.00
Good (≥70)	0.45 (0.30–0.67)	0.43 (0.29–0.64)	0.49 (0.33–0.72)	0.49 (0.25–0.97)	0.29 (0.16–0.51)	0.20 (0.10–0.43)
Mediterranean Diet Score	0.84 (0.73–0.97)	0.87 (0.74–1.04)	0.89 (0.77–1.03)	0.97 (0.82–1.16)	0.73 (0.56–0.94)	0.77 (0.57–1.04)
Non-adherers (0–4)	1.00	1.00	1.00	1.00	1.00	1.00
Adherers (5–9)	0.52 (0.33–0.82)	0.67 (0.41–1.11)	0.47 (0.29–0.76)	0.78 (0.47–1.32)	0.63 (0.33–1.21)	0.49 (0.18–1.37)

<sup>1</sup> Healthy Eating Index: Model 1: Adjusted for age at completion of the NHANES III questionnaire (continuous, years) and race/ethnicity (non-Hispanic white, non-Hispanic black, Mexican-American, other); Model 2: Additionally, adjusted for time between diagnosis and completion of the NHANES III questionnaire (continuous, years), body mass index (continuous, kg/m<sup>2</sup>), marital status (married/living together, never married/widowed, divorced/separated, unknown), socioeconomic status (poor/near poor, middle income, higher income, unknown), history of menopausal hormone therapy use (never user or no information, ever user), smoking status (never, former, or current smoker), self-reported prevalent chronic diseases at baseline (type 2 diabetes, heart attack, congestive heart failure or stroke; yes, no, or unknown), alcohol consumption (continuous, times per month), daily energy consumption (continuous, kcal/d) and moderate to vigorous physical activity (continuous, times of moderate to vigorous physical activity per week). Mediterranean Diet Score: Model 1: Adjusted for age at completion of the NHANES III questionnaire (continuous, years) and race/ethnicity (non-Hispanic white, non-Hispanic black, Mexican-American, other); Model 2: Additionally, adjusted for time between diagnosis and completion of the NHANES III questionnaire (continuous, years), body mass index (continuous, kg/m<sup>2</sup>), marital status (married/living together, never married/widowed, divorced/separated, unknown), socioeconomic status (poor/near poor, middle income, higher income, unknown), history of menopausal hormone therapy use (never user or no information, ever user), smoking status (never, former, or current smoker), self-reported prevalent chronic diseases at baseline (type 2 diabetes, heart attack, congestive heart failure or stroke; yes, no, or unknown), daily energy consumption (continuous, kcal/d) and moderate to vigorous physical activity (continuous, times of moderate to vigorous physical activity per week).



#### 4. Discussion

In our study of female cancer survivors, post-diagnostic diet quality, as assessed by the HEI, was associated with reduced mortality. In subgroup analyses, the statistically significant inverse association was evident in both breast and gynecological cancer survivors. For the MDS, a non-statistically significant inverse association was observed overall and in the subgroups.

Even though statistically significant associations were observed between the HEI and mortality, the association between the MDS and mortality was not statistically significant. A possible explanation for the lack of statistically significant associations is that very few participants in our study had high MDS. The MDS was based on the diet observed in European populations living around the Mediterranean region. On the other hand, the HEI was specifically designed to capture the diet quality of the US population (e.g., the first five components reflect conformity to the Food Pyramid Guide for the US population). This could suggest that the dietary pattern adopted by the US population differs from the Mediterranean diet and that the MDS may not be able to accurately capture the diet of this population. Given the multiple health benefits associated with the Mediterranean diet, several interventions were proposed in order to facilitate the adoption of the Mediterranean diet by the US population [27].

An inverse association between higher-quality diets (captured either by the HEI or the MDS) and mortality risk in disease-free populations was suggested previously [6,8,11,28,29]. Additionally, several studies have reported better prognosis in cancer survivors [30] and/or specifically female cancer survivors who adhered to a higher quality diet after diagnosis [4,10,31,32]. However, there are also studies suggesting that post-diagnostic diet and/or post-diagnostic diet quality are not associated with survival [5,19,33]. Our findings on the HEI are in line with the former studies and provide additional evidence in favor of a higher quality post-diagnostic diet as a modifiable risk factor.

A mechanism potentially explaining our findings could be mediated through inflammation. Breast cancer survivors who adopted higher quality diets post-diagnosis exhibited lower C-reactive protein (CRP) levels, compared to those with lower quality diets [34]. A positive association between CRP levels and mortality in disease-free individuals [35], as well as in cancer patients [36] is reported in the literature. Furthermore, improved survival after invasive breast cancer diagnosis was reported in women consuming diets with low inflammatory potential [37]. Diet with high inflammatory potential was previously associated with higher all-cause mortality risk in participants of the NHANES III [38]. Given that the diet corresponding to higher HEI score is considered a diet with low inflammatory potential, the inverse association we observed could be mediated through lower CRP concentrations. Another possible mechanism explaining our findings could be through meeting the micronutrient and mineral dietary requirements. Lower quality diets may be insufficient/deficient in various protective micronutrients and minerals and/or be abundant in potentially harmful agents. Long-term low micronutrient and mineral intake could be associated with worse health outcomes, including mortality [39,40].

The lifestyle differences and the lack of diet quality differences between female NHANES III participants without a self-reported cancer diagnosis and the female breast or gynecological cancer survivors included in this study (Table 2) could indicate that other lifestyle modifications, e.g., increasing physical activity or quitting smoking, may be easier to implement and to sustain over time compared to dietary modifications. Thus, further guidance and support may be necessary for female cancer survivors to adhere to a healthier, higher-quality diet post-diagnosis.

Our study had several strengths. The prospective study design and the long follow-up time allowed us to establish a clear period between disease onset, post-diagnostic dietary habits, and mortality. Additionally, the detailed information collection at baseline allowed us to adjust for various lifestyle factors that may influence mortality.

However, this study also has several limitations. The small sample size of female breast or gynecological cancer survivors may not have allowed the detection of a significant association between the MDS, and mortality. Additionally, our results were based on only one assessment of the dietary



habits and may not reflect the true or long-term habits of individual participants. However, both diet quality indices were validated in previous studies and seem to adequately capture peoples' dietary habits. As in any observational study, we cannot exclude the possibility of residual confounding. We cannot exclude the possibility that the female cancer survivors that participated in the NHANES III and in our study could have been healthier than female cancer survivors that did not participate in the NHANES III. Due to the small number of gynecological cancer survivors we were not able to perform analyses separately for every cancer type. It is plausible that some cancer cases (e.g., ovarian) were detected early since otherwise, patients' survival is very low. Finally, since information regarding disease severity or treatment was not available, we were not able to account for them in our analyses. Yet, given the long time between cancer diagnosis and mortality, cancer treatment and severity are unlikely to severely affect our results. Future projects should aim to capture this information and include them in the analyses as covariates.

In conclusion, higher post-diagnostic diet quality, as assessed by the HEI, was inversely associated with mortality in female cancer survivors, suggesting a protective effect when adhering to the diet captured by the HEI. Additional studies are required in order to verify our findings and to investigate underlying mechanisms of the mortality-adherence association.

**Disclaimer:** All analyses, interpretations, or conclusions reached are credited to the authors of this manuscript and not to the NCHS, which is responsible only for the initial data.

**Author Contributions:** Conceptualization, N.K. and S.R.; Methodology, N.K. and S.R.; Formal analysis, N.K.; Writing—original draft preparation, N.K.; Writing—review and editing, N.K., G.P., D.F. and S.R.; Supervision, S.R.; Funding acquisition, S.R.

**Funding:** This research was funded by Krebsliga Schweiz, grant number “KFS-4114-02-2017”.

**Conflicts of Interest:** The authors declare no conflict of interest. The funders had no role in the design of the study; in the collection, analyses, or interpretation of data; in the writing of the manuscript, or in the decision to publish the results.

## References

1. Ferlay, J.; Shin, H.; Bray, F.; Forman, D.; Mathers, C.; Parkin, D. *Globocan, Cancer Incidence and Mortality Worldwide: Iarc Cancerbase no. 10* [Internet]; International Agency for Research on Cancer: Lyon, France, 2010. Available online: [globocan.iarc.fr](http://globocan.iarc.fr) (accessed on 15 December 2017).
2. World Cancer Research Fund/American Institute for Cancer Research. Continuous Update Project Expert Report 2018. Diet, Nutrition, Physical Activity and Breast Cancer Survivors. Available online: [dietandcancerreport.org](http://dietandcancerreport.org) (accessed on 15 December 2017).
3. Pierce, J.P.; Stefanick, M.L.; Flatt, S.W.; Natarajan, L.; Sternfeld, B.; Madlensky, L.; Al-Delaimy, W.K.; Thomson, C.A.; Kealey, S.; Hajek, R. Greater survival after breast cancer in physically active women with high vegetable-fruit intake regardless of obesity. *J. Clin. Oncol.* **2007**, *25*, 2345–2351. [CrossRef] [PubMed]
4. Inoue-Choi, M.; Robien, K.; Lazovich, D. Adherence to the wcrf/aicr guidelines for cancer prevention is associated with lower mortality among older female cancer survivors. *Cancer Epidemiol. Biomark. Prev.* **2013**, *22*, 792–802. [CrossRef] [PubMed]
5. Pierce, J.P.; Natarajan, L.; Caan, B.J.; Parker, B.A.; Greenberg, E.R.; Flatt, S.W.; Rock, C.L.; Kealey, S.; Al-Delaimy, W.K.; Bardwell, W.A.; et al. Influence of a diet very high in vegetables, fruit, and fiber and low in fat on prognosis following treatment for breast cancer: The women's healthy eating and living (whel) randomized trial. *JAMA* **2007**, *298*, 289–298. [CrossRef] [PubMed]
6. Rathod, A.D.; Bharadwaj, A.S.; Badheka, A.O.; Kizilbash, M.; Afonso, L. Healthy eating index and mortality in a nationally representative elderly cohort. *Arch. Intern. Med.* **2012**, *172*, 275–277. [CrossRef] [PubMed]
7. Reedy, J.; Krebs-Smith, S.M.; Miller, P.E.; Liese, A.D.; Kahle, L.L.; Park, Y.; Subar, A.F. Higher diet quality is associated with decreased risk of all-cause, cardiovascular disease, and cancer mortality among older adults. *J. Nutr.* **2014**, *144*, 881–889. [CrossRef] [PubMed]

8. George, S.M.; Ballard-Barbash, R.; Manson, J.E.; Reedy, J.; Shikany, J.M.; Subar, A.F.; Tinker, L.F.; Vitolins, M.; Neuhouser, M.L. Comparing indices of diet quality with chronic disease mortality risk in postmenopausal women in the women's health initiative observational study: Evidence to inform national dietary guidance. *Am. J. Epidemiol.* **2014**, *180*, 616–625. [CrossRef] [PubMed]
9. Onvani, S.; Haghighatdoost, F.; Surkan, P.J.; Larijani, B.; Azadbakht, L. Adherence to the healthy eating index and alternative healthy eating index dietary patterns and mortality from all causes, cardiovascular disease and cancer: A meta-analysis of observational studies. *J. Hum. Nutr. Diet.* **2017**, *30*, 216–226. [CrossRef]
10. George, S.M.; Ballard-Barbash, R.; Shikany, J.M.; Caan, B.J.; Freudenheim, J.L.; Kroenke, C.H.; Vitolins, M.Z.; Beresford, S.A.; Neuhouser, M.L. Better postdiagnosis diet quality is associated with reduced risk of death among postmenopausal women with invasive breast cancer in the women's health initiative. *Cancer Epidemiol. Biomark. Prev.* **2014**, *23*, 575–583. [CrossRef]
11. Knoops, K.T.; de Groot, L.C.; Kromhout, D.; Perrin, A.E.; Moreiras-Varela, O.; Menotti, A.; van Staveren, W.A. Mediterranean diet, lifestyle factors, and 10-year mortality in elderly european men and women: The hale project. *JAMA* **2004**, *292*, 1433–1439. [CrossRef]
12. Osler, M.; Schroll, M. Diet and mortality in a cohort of elderly people in a north european community. *Int. J. Epidemiol.* **1997**, *26*, 155–159. [CrossRef]
13. Trichopoulou, A.; Kouris-Blazos, A.; Wahlqvist, M.L.; Gnardellis, C.; Lagiou, P.; Polychronopoulos, E.; Vassilakou, T.; Lipworth, L.; Trichopoulos, D. Diet and overall survival in elderly people. *BMJ* **1995**, *311*, 1457–1460. [CrossRef] [PubMed]
14. Trichopoulou, A.; Costacou, T.; Bamia, C.; Trichopoulos, D. Adherence to a mediterranean diet and survival in a greek population. *N. Engl. J. Med.* **2003**, *348*, 2599–2608. [CrossRef] [PubMed]
15. Trichopoulou, A.; Bamia, C.; Trichopoulos, D. Mediterranean diet and survival among patients with coronary heart disease in greece. *Arch. Intern. Med.* **2005**, *165*, 929–935. [CrossRef] [PubMed]
16. Lopez-Garcia, E.; Rodriguez-Artalejo, F.; Li, T.Y.; Fung, T.T.; Li, S.; Willett, W.C.; Rimm, E.B.; Hu, F.B. The mediterranean-style dietary pattern and mortality among men and women with cardiovascular disease. *Am. J. Clin. Nutr.* **2013**, *99*, 172–180. [CrossRef]
17. Bonaccio, M.; Di Castelnuovo, A.; Costanzo, S.; Persichillo, M.; De Curtis, A.; Donati, M.B.; de Gaetano, G.; Iacoviello, L.; Investigators, M.-S. Adherence to the traditional mediterranean diet and mortality in subjects with diabetes. Prospective results from the moli-sani study. *Eur. J. Prev. Cardiol.* **2016**, *23*, 400–407. [CrossRef]
18. Schwingshackl, L.; Schwedhelm, C.; Galbete, C.; Hoffmann, G. Adherence to mediterranean diet and risk of cancer: An updated systematic review and meta-analysis. *Nutrients* **2017**, *9*, 1063. [CrossRef]
19. Kim, E.H.; Willett, W.C.; Fung, T.; Rosner, B.; Holmes, M.D. Diet quality indices and postmenopausal breast cancer survival. *Nutr. Cancer* **2011**, *63*, 381–388. [CrossRef]
20. US Department of Health and Human Services, P.H.S., Centers for Disease Control and Prevention. Plan and operation of the third national health and nutrition examination survey, 1988–1994. In *Vital Health Stat 1994*, 1; US Department of Health and Human Services: Hyattsville, MD, USA, 1994.
21. NHANES III (1988–1994). Available online: <https://wwwn.cdc.gov/nchs/nhanes/nhanes3/Default.aspx> (accessed on 15 December 2017).
22. NCHS Data Linkage–Mortality Data–Methods. Available online: <http://cdc.gov/nchs/data-linkage/mortality-methods.htm> (accessed on 15 December 2017).
23. National Center for Health Statistics. *National Death Index User's Guide*; National Center for Health Statistics: Hyattsville, MD, USA, 2013.
24. Calle, E.E.; Terrell, D.D. Utility of the national death index for ascertainment of mortality among cancer prevention study ii participants. *Am. J. Epidemiol.* **1993**, *137*, 235–241. [CrossRef]
25. Rich-Edwards, J.W.; Corsano, K.A.; Stampfer, M.J. Test of the national death index and equifax nationwide death search. *Am. J. Epidemiol.* **1994**, *140*, 1016–1019. [CrossRef]
26. Suresh, S.; Sabanayagam, C.; Shankar, A. Socioeconomic status, self-rated health, and mortality in a multiethnic sample of us adults. *J. Epidemiol.* **2011**, *21*, 337–345. [CrossRef]
27. Martínez-González, M.Á.; Hershey, M.S.; Zazpe, I.; Trichopoulou, A. Transferability of the mediterranean diet to non-mediterranean countries. What is and what is not the mediterranean diet. *Nutrients* **2017**, *9*, 1226. [CrossRef] [PubMed]

28. Akbaraly, T.N.; Ferrie, J.E.; Berr, C.; Brunner, E.J.; Head, J.; Marmot, M.G.; Singh-Manoux, A.; Ritchie, K.; Shipley, M.J.; Kivimaki, M. Alternative healthy eating index and mortality over 18 years of follow-up: Results from the whitehall ii cohort. *Am. J. Clin. Nutr.* **2011**, *94*, 247–253. [[CrossRef](#)] [[PubMed](#)]
29. Mitrou, P.N.; Kipnis, V.; Thiebaut, A.C.; Reedy, J.; Subar, A.F.; Wirfalt, E.; Flood, A.; Mouw, T.; Hollenbeck, A.R.; Leitzmann, M.F.; et al. Mediterranean dietary pattern and prediction of all-cause mortality in a us population: Results from the nih-aarp diet and health study. *Arch. Intern. Med.* **2007**, *167*, 2461–2468. [[CrossRef](#)] [[PubMed](#)]
30. Schwedhelm, C.; Boeing, H.; Hoffmann, G.; Aleksandrova, K.; Schwingshackl, L. Effect of diet on mortality and cancer recurrence among cancer survivors: A systematic review and meta-analysis of cohort studies. *Nutr. Rev.* **2016**, *74*, 737–748. [[CrossRef](#)] [[PubMed](#)]
31. Jochems, S.H.J.; Van Osch, F.H.M.; Bryan, R.T.; Wesselius, A.; van Schooten, F.J.; Cheng, K.K.; Zeegers, M.P. Impact of dietary patterns and the main food groups on mortality and recurrence in cancer survivors: A systematic review of current epidemiological literature. *BMJ Open* **2018**, *8*, 14530. [[CrossRef](#)] [[PubMed](#)]
32. Terranova, C.O.; Protani, M.M.; Reeves, M.M. Overall dietary intake and prognosis after breast cancer: A systematic review. *Nutr. Cancer* **2018**, *70*, 153–163. [[CrossRef](#)] [[PubMed](#)]
33. Izano, M.A.; Fung, T.T.; Chiuve, S.S.; Hu, F.B.; Holmes, M.D. Are diet quality scores after breast cancer diagnosis associated with improved breast cancer survival? *Nutr. Cancer* **2013**, *65*, 820–826. [[CrossRef](#)]
34. George, S.M.; Neuhouser, M.L.; Mayne, S.T.; Irwin, M.L.; Albanes, D.; Gail, M.H.; Alfano, C.M.; Bernstein, L.; McTiernan, A.; Reedy, J.; et al. Postdiagnosis diet quality is inversely related to a biomarker of inflammation among breast cancer survivors. *Cancer Epidemiol. Prev. Biomark.* **2010**, *19*, 2220–2228. [[CrossRef](#)]
35. Reuben, D.B.; Cheh, A.I.; Harris, T.B.; Ferrucci, L.; Rowe, J.W.; Tracy, R.P.; Seeman, T.E. Peripheral blood markers of inflammation predict mortality and functional decline in high-functioning community-dwelling older persons. *J. Am. Geriatr. Soc.* **2002**, *50*, 638–644. [[CrossRef](#)]
36. McMillan, D.C. The systemic inflammation-based glasgow prognostic score: A decade of experience in patients with cancer. *Cancer Treat. Rev.* **2013**, *39*, 534–540. [[CrossRef](#)]
37. Zheng, J.; Tabung, F.K.; Zhang, J.; Liese, A.D.; Shivappa, N.; Ockene, J.K.; Caan, B.; Kroenke, C.H.; Hebert, J.R.; Steck, S.E. Association between post-cancer diagnosis dietary inflammatory potential and mortality among invasive breast cancer survivors in the women’s health initiative. *Cancer Epidemiol. Prev. Biomark.* **2018**, *27*, 454–463. [[CrossRef](#)] [[PubMed](#)]
38. Shivappa, N.; Steck, S.E.; Hussey, J.R.; Ma, Y.; Hebert, J.R. Inflammatory potential of diet and all-cause, cardiovascular, and cancer mortality in national health and nutrition examination survey iii study. *Eur. J. Nutr.* **2017**, *56*, 683–692. [[CrossRef](#)] [[PubMed](#)]
39. Bell, G.A.; Kantor, E.D.; Lampe, J.W.; Kristal, A.R.; Heckbert, S.R.; White, E. Intake of long-chain omega-3 fatty acids from diet and supplements in relation to mortality. *Am. J. Epidemiol.* **2014**, *179*, 710–720. [[CrossRef](#)] [[PubMed](#)]
40. Micha, R.; Penalvo, J.L.; Cudhea, F.; Imamura, F.; Rehm, C.D.; Mozaffarian, D. Association between dietary factors and mortality from heart disease, stroke, and type 2 diabetes in the united states. *JAMA* **2017**, *317*, 912–924. [[CrossRef](#)] [[PubMed](#)]

